→ USPTO

11/13/2008 12:50 FAX 6099219535

- 1. (CURRENTLY AMENDED) A catalyst comprising a composite complexmade of one or more metal species having 10 to 50,000 atoms and an organicmatter bonded to the metal species, the composite complex carried on a calcined-The method of claim 14 wherein the porous carrier made of comprises one or more metal oxides or carbon.
- 2. (CURRENTLY AMENDED) The entalyst method according to claim 14 1, wherein the composite complex is reduced before or after calcining the carrier having carried the composite complex thereon.
- 3. (CURRENTLY AMENDED) The catalyst method according to claim 14 1, wherein the composite complex is soluble in water.
- 4. (CURRENTLY AMENDED) The eatalyst method according to claim 14 1, wherein the organic matter composing the composite complex has at least one of nitrogen, a carboxyl group, a carbonyl group and an alcohol group.
- 5. (CANCELED)
- 6. (CANCELED)
- 7. (CURRENTLY AMENDED) The estalyst method according to claim 14 1, wherein the one or more metal species comprises ions comprise of one or more of platinum, palladium, rhodium, ruthenium, silver, gold and iridium.

- 8. (CURRENTLY AMENDED) The eatalyst method according to claim 14 1, wherein the porous carrier comprises at least one of ceria, ceria-zirconia, ceria-zirconia-yttria, alumina, silica, titania and zirconia.
- 9. (CURRENTLY AMENDED) The catalyst method according to claim 14 1, wherein the porous carrier comprises at least one of aluminum, zirconium, silicon, titanium, lanthanum, cerium, neodymium and yttrium.
- 10. (CURRENTLY AMENDED) The eatalyst method according to claim 14 1, further comprising at least one of an alkali metal element, an alkaline earth metal element and a rare earth metal element carried thereon.
- 11. (CURRENTLY AMENDED) The eatalyst method according to claim 10, wherein the alkali metal element is lithium, sodium, potassium, rubidium or cesium.
- 12. (CURRENTLY AMENDED) The eatalyst method according to claim 10, wherein the alkaline earth metal element is magnesium, calcium, strontium or barium.
- 13. (CURRENTLY AMENDED) The eatelyst method according to claim 10, wherein the rare earth metal element is lanthanum or cerium.
- 14. (ORIGINAL) A method for producing a catalyst comprising the steps of: producing a metal salt solution containing salts of one or more metals; dispersing the metal salt solution, an organic matter and a porous carrier made of one or more metal oxides in a solvent to form a composite complex comprising one or

more metal ions having 10 to 50,000 atoms and the organic matter bonded to the metal ions, and to simultaneously make the composite complex carried on the porous carrier; and calcining the carrier having the composite complex carried thereon.

- 15. (ORIGINAL) The method for producing a catalyst according to claim 14, further comprising a step of reducing the metal ions on the porous carrier by reducing the carrier after the step of making the composite complex carried on the carrier and before or after calcining the carrier, and then calcining the carrier after the reduction step.
- 16. (PREVIOUSLY PRESENTED) The method for producing a catalyst according to claim 14, wherein the organic matter has at least one of nitrogen, a carboxyl group, a carboxyl group and an alcohol group.
- 17. (PREVIOUSLY PRESENTED) The method for producing a catalyst according to claim 16, wherein the organic matter comprises at least one of polyacrylic acid, polymethacrylic acid, polyethyleneimine, polyallylamine, polyvinylpyrrolidone, polyvinyl alcohol, poly(N-carboxymethyl)ethyleneimine, ,N-d icarboxymethyl)arylamine, or a copolymer containing one or more thereof.
- 18. (PREVIOUSLY PRESENTED) The method for producing a catalyst according to claim 16, wherein the organic matter is comprises polyamino acid or polysaccharide.
- 19. (CURRENTLY AMENDED) The catalyst method according to claim 14 2, wherein the composite complex is soluble in water.

11/13/2008 12:50 FAX 6099219535

20. (CURRENTLY AMENDED) The eatalyst method according to claim 14 2, wherein the organic matter composing the composite complex has at least one of nitrogen, a carboxyl group, a carbonyl group and an alcohol group.